


This listing of claims will replace all prior versions, and listings, of claims in the application:

**Listing of Claims:**

1. (Previously Presented): A cationic dye of formula I




wherein

CAT<sup>+</sup> is a cation selected from azine, xanthene, polymethine, styryl, azo, tetrazolium, pyrylium, benzopyrylium, thiopyrylium, benzothiopyrylium, thiazine, oxazine, triarylmethane, diarylmethane, acridine, quinoline, isoquinoline, and quaternized azafluorenone dyes,

Y<sup>-</sup> is an anion selected from CAB<sup>-</sup>, FAP<sup>-</sup>, FAB<sup>-</sup>, and Im<sup>-</sup>,

CAB<sup>-</sup> conforms to formula (II-1)




y1 is 1, 2, 3 or 4,

x1 is 0, 1, 2 or 3,

R<sup>0</sup> is alkyl, aryl, fluorinated alkyl, fluorinated aryl, cycloalkyl or alkylaryl, with the condition that R<sup>0</sup> may be hydrogen if y1 is >2,

FAP<sup>-</sup> conforms to formula (II-2)




p2 is 1 to 20,

m2 is 0, 1, 2 or 3,

y2 is 1, 2, 3 or 4,

FAB<sup>-</sup> conforms to formula (II-3)




p3 is 1 to 20,

m3 is 0, 1, 2 or 3,

y3 is 1, 2, 3 or 4,

**MERCK-3134**

Im<sup>-</sup> conforms to formula (II-4)



X is carbon or sulfur,

p<sub>4</sub> is 0 to 20 and  $0 \leq m_4 \leq 2p_4+1$ ,

q is 0 to 20 and  $0 \leq k \leq 2q+1$ ,

y<sub>4</sub> is 1 or 2,

m<sub>4</sub> is 0 if p<sub>4</sub> is 0,

k is 0 if q is 0, and

the carbon atoms of the alkyl chain of the formula II-4 may be bonded to one another by single bonds, where the resultant alkylene chain may in turn be partially or fully substituted by F;

with the provisos that:

if X is sulfur, y<sub>4</sub> is 2, and if X is carbon, y<sub>4</sub> is 1 and p<sub>4</sub> or q ≥ 1, and

3,3'-diethoxyethyl-2,2'-thiadibenzocyanine trifluoromethyltrifluoroborate is excluded.

2. (Withdrawn): A dye according to Claim 1, wherein CAT<sup>+</sup> is a cation of an azine dye.

3. (Withdrawn): A dye according to Claim 1, wherein CAT<sup>+</sup> is a cation of a xanthene dye.

4. (Previously Presented): A dye according to Claim 1, wherein CAT<sup>+</sup> is a cation of a polymethine dye.

5. (Withdrawn): A dye according to Claim 1, wherein CAT<sup>+</sup> is a cation of a styryl dye.

6. (Withdrawn): A dye ~~Dyes~~ according to Claim 1, wherein CAT<sup>+</sup> is a cation of an azo dye.

**MERCK-3134**

7. (Withdrawn): A dye according to Claim 1, wherein  $CAT^+$  is a cation of a tetrazolium dye.

8. (Withdrawn): A dye according to Claim 1, wherein  $CAT^+$  is a cation of a pyrylium dye.

9. (Withdrawn): A dye according to Claim 1, wherein  $CAT^+$  is a cation of a benzopyrylium dye.

10. (Withdrawn): A dye according to Claim 1, wherein  $CAT^+$  is a cation of a thiopyrylium dye.

11. (Withdrawn): A dye according to Claim 1, wherein  $CAT^+$  is a cation of a benzothiopyrylium dye.

12. (Withdrawn): A dye according to Claim 1, wherein  $CAT^+$  is a cation of a thiazine dye.

13. (Withdrawn): A dye according to Claim 1, wherein  $CAT^+$  is a cation of an oxazine dye.


14. (Withdrawn): A dye ~~Dyes~~ according to Claim 1, wherein  $CAT^+$  is a cation of a triarylmethane dye.

15. (Withdrawn; Currently Amended): A dye according to Claim 1, wherein  $CAT^+$  is a cation of a diarylmethane dye.

16. (Withdrawn): A dye according to Claim 1, wherein  $CAT^+$  is a cation of an acridine dye.

17. (Withdrawn): A dye according to Claim 1, wherein  $CAT^+$  is a cation of a quinoline dye.
18. (Withdrawn): A dye according to Claim 1, wherein  $CAT^+$  is a cation of an isoquinoline dye.
19. (Withdrawn): A dye according to Claim 1, wherein  $CAT^+$  is a cation of a quaternary azafluorenone dye.
20. (Previously Presented): A dye according to Claim 4, wherein  $CAT^+$  is a cation of a cyanine dye.
21. (Previously Presented): A dye according to Claim 4, wherein  $CAT^+$  is a cation of a carbocyanine dye.
22. (Previously Presented): A dye according to Claim 4, wherein  $CAT^+$  is a cation of an azacarbocyanine dye.
23. (Previously Presented): A dye according to Claim 4, wherein  $CAT^+$  is a cation of a diazacarbocyanine dye.
24. (Previously Presented): A dye according to Claim 4, wherein  $CAT^+$  is a cation of a triazacarbocyanine dye.
25. (Previously Presented): A dye according to Claim 4, wherein  $CAT^+$  is a cation of a hemicyanine dye.
26. (Previously Presented): A dye according to Claim 4, wherein  $CAT^+$  is a cation of a diazahemicyanine dye.
27. (Withdrawn): A dye according to claim 1, wherein  $Y^-$  is a cyanoborate of

formula II-1




wherein

$y1$  is 1, 2, 3 or 4,

$x1$  is 0, 1, 2 or 3 and

$\text{R}^0$  is alkyl, aryl, fluorinated alkyl, fluorinated aryl, cycloalkyl or alkylaryl, with the condition that  $\text{R}^0$  may be hydrogen if  $y1$  is  $>2$ .

28. (Previously Presented): A dye according to claim 1, wherein  $\text{Y}^-$  is a fluoroalkylphosphate of the formula II-2




wherein

$p2$  is 1 to 20,

$m2$  is 0, 1, 2 or 3 and

$y2$  is 1, 2, 3 or 4 .

29. (Withdrawn): A dye according to claim 1, wherein  $\text{Y}^-$  is a fluoroalkylborate of formula II-3




wherein

$p3$  is 1 to 20,

$m3$  is 0, 1, 2 or 3 and

$y3$  is 1, 2, 3 or 4.

30. (Withdrawn): A dye according to claim 1, wherein  $\text{Y}^-$  is an imide of the formula II-4



wherein

$\text{X}$  is carbon or sulfur,

$p4$  is 0 to 20 and  $0 \leq m4 \leq 2p4+1$ ,

$q$  is 0 to 20 and  $0 \leq k \leq 2q+1$ ,

**MERCK-3134**

y<sub>4</sub> is 1 or 2,  
 m<sub>4</sub> is 0 if p<sub>4</sub> is 0, and  
 k is 0 if q is 0,

with the proviso that

if X is sulfur, y<sub>4</sub> is 2, and if X is carbon, y<sub>4</sub> is 1 and p<sub>4</sub> or q ≥ 1.


31. (Withdrawn): A process for the preparation of a cationic dye according to claim 1, said process comprising:

reacting a compound of formula XXI



wherein A<sup>-</sup> is Cl<sup>-</sup>, Br<sup>-</sup>, I<sup>-</sup>, BF<sub>4</sub><sup>-</sup>, PF<sub>6</sub><sup>-</sup>, ClO<sub>4</sub><sup>-</sup>, sulfate, tosylate, hydrosulfate, triflate, trifluoroacetate, acetate or oxalate,


with a compound of formula XXII



wherein Y<sup>-</sup> is CAB<sup>-</sup>, FAP<sup>-</sup>, FAB<sup>-</sup> or Im<sup>-</sup>, and

E<sup>+</sup> is a cation selected from cations of the alkali metals, alkaline earth metals or of a metal from group 11 and 12, ammonium, alkylammonium containing C<sub>1</sub>-C<sub>4</sub>-alkyl, phosphonium, alkylphosphonium containing C<sub>1</sub>-C<sub>4</sub>-alkyl, and ox guanidinium.

32. (Withdrawn): A process for the preparation of carbocyanine dye according to Claim 21, where the carbocyanine dye conforms to formula XXIII



wherein

n is 0, 1, 2, 3, 4 or 5,


R in each case, independently of one another, is alkyl, alkenyl, cycloalkyl, aryl or heteroaryl,

R<sup>1</sup> in each case, independently of one another, is H, Cl, Br, I, alkyl, partially or fully chlorinated alkyl, alkenyl, cycloalkyl, aryl, heteroaryl, Oalkyl, Oaryl, Salkyl, Saryl,

**MERCK-3134**

NHalkyl, N(alkyl)<sub>2</sub>, C(O)H, C(O)alkyl, C(O)aryl, CN, N=N-aryl, P(aryl)<sub>2</sub>, NHC(O)alkyl or NHC(O)aryl and

the ring system, represented by




is a nitrogen-containing unsaturated mono-, bi- or tricyclic heterocycle having 5 to 13 ring members, which optionally contains 1, 2 or 3 N and/or 1 or 2 S or O atoms and the heterocyclic radical is optionally mono- or polysubstituted by Z,

Z is hydrogen, alkyl, NO<sub>2</sub>, F, Cl, Br, I, OH, COOH, Oalkyl, SCN, SCF<sub>3</sub>, COOalkyl, CH<sub>2</sub>-COOalkyl, NH<sub>2</sub>, NHalkyl or N(alkyl)<sub>2</sub>

Y<sup>-</sup> is an anion selected from CAB<sup>-</sup>, FAP<sup>-</sup>, FAB and Im<sup>-</sup>,

CAB<sup>-</sup> conforms to formula (II-1)




y<sub>1</sub> is 1, 2, 3 or 4,

x<sub>1</sub> is 0, 1, 2 or 3,

R<sup>0</sup> is alkyl, aryl, fluorinated alkyl, fluorinated aryl, cycloalkyl or alkylaryl, with the condition that R<sup>0</sup> may be hydrogen if y<sub>1</sub> is >2,

FAP<sup>-</sup> conforms to formula (II-2)




p<sub>2</sub> is 1 to 20,

m<sub>2</sub> is 0, 1, 2 or 3,

y<sub>2</sub> is 1, 2, 3 or 4,

FAB<sup>-</sup> conforms to formula (II-3)




p<sub>3</sub> is 1 to 20,

m<sub>3</sub> is 0, 1, 2 or 3,

y<sub>3</sub> is 1, 2, 3 or 4,

Im<sup>-</sup> conforms to formula (II-4)



X is carbon or sulfur,

p<sub>4</sub> is 0 to 20 and  $0 \leq m_4 \leq 2p_4+1$ ,

q is 0 to 20 and  $0 \leq k \leq 2q+1$ ,

y<sub>4</sub> is 1 or 2,

m<sub>4</sub> is 0 if p<sub>4</sub> is 0,


k is 0 if q is 0, and

the carbon atoms of the alkyl chain of the formula II-4 may be bonded to one another by single bonds, where the resultant alkylene chain may in turn be partially or fully substituted by F;

with the proviso that

if X is sulfur, y<sub>4</sub> is 2, and if X is carbon, y<sub>4</sub> is 1 and p<sub>4</sub> or q ≥ 1,

said process comprising utilizing a compound of formula XXIV




where the ring system, R, R<sup>1</sup> and Y<sup>-</sup> have one of the meanings indicated in the case of formula XXIII, and

n is 0, 1, 2, 3 or 4 and

G is hydrogen, alkyl, alkenyl, aryl, heteroaryl, N=C(R)<sub>2</sub>, CONHaryl, C(O)aryl or CONHalkyl.

33. (Withdrawn): A compound according to formula XXIV



where

**MERCK-3134**




n is 0, 1, 2, 3 or 4,

G is hydrogen, alkyl, alkenyl, aryl, heteroaryl, N=C(R)<sub>2</sub>, CONHaryl, C(O)aryl or CONHalkyl,

R is alkyl, alkenyl, cycloalkyl, aryl or heteroaryl,

R<sup>1</sup> is in each case, independently of one another, H, Cl, Br, I, alkyl, partially or fully chlorinated alkyl, alkenyl, cycloalkyl, aryl, heteroaryl, Oalkyl, Oaryl, Salkyl, Saryl, NHalkyl, N(alkyl)<sub>2</sub>, C(O)H, C(O)alkyl, C(O)aryl, CN, N=N-aryl, P(aryl)<sub>2</sub>, NHC(O)alkyl or NHC(O)aryl, and

the ring system, represented by




is a nitrogen-containing unsaturated mono-, bi- or tricyclic heterocycle having 5 to 13 ring members, optionally containing 1, 2 or 3 N and/or 1 or 2 S or O atoms and in which the heterocyclic radical is optionally mono- or polysubstituted by Z,

Z is hydrogen, alkyl, NO<sub>2</sub>, F, Cl, Br, I, OH, COOH, Oalkyl, SCN, SCF<sub>3</sub>, COOalkyl, CH<sub>2</sub>-COOalkyl, NH<sub>2</sub>, NHalkyl or N(alkyl)<sub>2</sub>,

Y<sup>-</sup> is an anion selected from CAB<sup>-</sup>, FAP<sup>-</sup>, FAB<sup>-</sup> and Im<sup>-</sup>,

CAB<sup>-</sup> conforms to formula (II-1)




y<sub>1</sub> is 1, 2, 3 or 4,

x<sub>1</sub> is 0, 1, 2 or 3,

R<sup>0</sup> is alkyl, aryl, fluorinated alkyl, fluorinated aryl, cycloalkyl or alkylaryl, with the condition that R<sup>0</sup> may be hydrogen if y<sub>1</sub> is >2,

FAP<sup>-</sup> conforms to formula (II-2)




p<sub>2</sub> is 1 to 20,

m<sub>2</sub> is 0, 1, 2 or 3,

y<sub>2</sub> is 1, 2, 3 or 4,

FAB<sup>-</sup> conforms to formula (II-3)




p<sub>3</sub> is 1 to 20,

m<sub>3</sub> is 0, 1, 2 or 3,

y<sub>3</sub> is 1, 2, 3 or 4,

Im<sup>-</sup> conforms to formula (II-4)



X is carbon or sulfur,

p<sub>4</sub> is 0 to 20 and  $0 \leq m_4 \leq 2p_4+1$ ,

q is 0 to 20 and  $0 \leq k \leq 2q+1$ ,

y<sub>4</sub> is 1 or 2,

m<sub>4</sub> is 0 if p<sub>4</sub> is 0, and

k is 0 if q is 0,

where the carbon atoms of the alkyl chain of the formula II-4 may be bonded to one another by single bonds, and the resultant alkylene chain may in turn be partially or fully substituted by F;


with the provisos that:

if X is sulfur, y<sub>4</sub> is 2, and

if X is carbon, y<sub>4</sub> is 1 and p<sub>4</sub> or q ≥ 1.

34. (Withdrawn): A process for the preparation of a compound according to Claim 33, said process comprising reacting

a compound of formula XXV




in which

A<sup>-</sup> is Cl<sup>-</sup>, Br<sup>-</sup>, I<sup>-</sup>, BF<sub>4</sub><sup>-</sup>, PF<sub>6</sub><sup>-</sup>, ClO<sub>4</sub><sup>-</sup>, sulfate, tosylate, hydrosulfate, triflate,

**MERCK-3134**

trifluoroacetate, acetate or oxalate,

the ring system, represented by



is a nitrogen-containing unsaturated mono-, bi- or tricyclic heterocycle having 5 to 13 ring members, which optionally further contains 1, 2 or 3 N and/or 1 or 2 S or O atoms, and in which the heterocyclic radical is optionally mono- or polysubstituted by Z,

Z is hydrogen, alkyl, NO<sub>2</sub>, F, Cl, Br, I, OH, COOH, Oalkyl, SCN, SCF<sub>3</sub>, COOalkyl, CH<sub>2</sub>-COOalkyl, NH<sub>2</sub>, NHalkyl, or N(alkyl)<sub>2</sub>,

n is 0, 1, 2, 3 or 4,

R is alkyl, alkenyl, cycloalkyl, aryl or heteroaryl,

R<sup>1</sup> is in each case, independently of one another, H, Cl, Br, I, alkyl, partially or fully chlorinated alkyl, alkenyl, cycloalkyl, aryl, heteroaryl, Oalkyl, Oaryl, Salkyl, Saryl, NHalkyl, N(alkyl)<sub>2</sub>, C(O)H, C(O)alkyl, C(O)aryl, CN, N=N-aryl, P(aryl)<sub>2</sub>, NHC(O)alkyl, or NHC(O)aryl, and

G is hydrogen, alkyl, alkenyl, aryl, heteroaryl, N=C(R)<sub>2</sub>, CONHaryl, C(O)aryl, or CONHalkyl,

with a compound of formula XXVI




in which

E<sup>+</sup> is a cation of the alkali metals, alkaline earth metals or of a metal from group 11 and 12, ammonium, alkylammonium containing C<sub>1</sub>-C<sub>4</sub>-alkyl, phosphonium, alkylphosphonium containing C<sub>1</sub>-C<sub>4</sub>-alkyl, or guanidinium,

Y<sup>-</sup> is an anion selected from CAB<sup>-</sup>, FAP<sup>-</sup>, FAB<sup>-</sup> and Im<sup>-</sup>,


CAB<sup>-</sup> conforms to formula (II-1)



**MERCK-3134**


y1 is 1, 2, 3 or 4,  
 x1 is 0, 1, 2 or 3,  
 R<sup>0</sup> is alkyl, aryl, fluorinated alkyl, fluorinated aryl, cycloalkyl or alkylaryl, with  
 the condition that R<sup>0</sup> may be hydrogen if y1 is >2,

FAP<sup>-</sup> conforms to formula (II-2)




p2 is 1 to 20,  
 m2 is 0, 1, 2 or 3,  
 y2 is 1, 2, 3 or 4,

FAB<sup>-</sup> conforms to formula (II-3)



p3 is 1 to 20,  
 m3 is 0, 1, 2 or 3,  
 y3 is 1, 2, 3 or 4,

Im<sup>-</sup> conforms to formula (II-4)



X is carbon or sulfur,  
 p4 is 0 to 20 and  $0 \leq m4 \leq 2p4+1$ ,  
 q is 0 to 20 and  $0 \leq k \leq 2q+1$ ,  
 y4 is 1 or 2,  
 m4 is 0 if p4 is 0, and  
 k is 0 if q is 0,

where the carbon atoms of the alkyl chain of the formula II-4 may be bonded to one another by single bonds, and the resultant alkylene chain may in turn be partially or fully substituted by F;


with the provisos that

if X is sulfur, y4 is 2, and if X is carbon, y4 is 1 and p4 or q ≥ 1.

35. (Withdrawn): A process for the preparation of a compound according to Claim 33, with the restriction that n in formula XXIV is 0, said process comprising:

**MERCK-3134**

reacting a compound of the formula XXVII




in which

G is hydrogen, alkyl, alkenyl, aryl, heteroaryl, N=C(R)<sub>2</sub>, CONHaryl, C(O)aryl, or CONHalkyl,

R is alkyl, alkenyl, cycloalkyl, aryl or heteroaryl,

the ring system, represented by



is a nitrogen-containing unsaturated mono-, bi- or tricyclic heterocycle having 5 to 13 ring members, which optionally further contains 1, 2 or 3 N and/or 1 or 2 S or O atoms, and in which the heterocyclic radical is optionally mono- or polysubstituted by Z,


Z is hydrogen, alkyl, NO<sub>2</sub>, F, Cl, Br, I, OH, COOH, Oalkyl, SCN, SCF<sub>3</sub>, COOalkyl, CH<sub>2</sub>-COOalkyl, NH<sub>2</sub>, NHalkyl, or N(alkyl)<sub>2</sub>,

with a compound HY,

where

Y<sup>-</sup> is an anion selected from FAP<sup>-</sup>, FAB<sup>-</sup> and Im<sup>-</sup>,

FAP<sup>-</sup> conforms to formula (II-2)




p<sub>2</sub> is 1 to 20,

m<sub>2</sub> is 0, 1, 2 or 3,

y<sub>2</sub> is 1, 2, 3 or 4,


FAB<sup>-</sup> conforms to formula (II-3)



**MERCK-3134**

p3 is 1 to 20,  
 m3 is 0, 1, 2 or 3,  
 y3 is 1, 2, 3 or 4,

Im<sup>-</sup> conforms to formula (II-4)




X is carbon or sulfur,  
 p4 is 0 to 20 and  $0 \leq m4 \leq 2p4+1$ ,  
 q is 0 to 20 and  $0 \leq k \leq 2q+1$ ,  
 y4 is 1 or 2,  
 m4 is 0 if p4 is 0, and  
 k is 0 if q is 0,

where the carbon atoms of the alkyl chain of the formula II-4 may be bonded to one another by single bonds, and the resultant alkylene chain may in turn be partially or fully substituted by F;

with the provisos that

if X is sulfur, y4 is 2, and if X is carbon, y4 is 1 and p4 or q  $\geq$  1.

36. (Withdrawn): A process for the preparation of an azo dyes according to Claim 6, wherein said azo dye conforms to formula XXVIII




where

R' and R'' are each aryl or heteroaryl and one of the two aromatic nuclei is positively charged,

Y<sup>-</sup> is an anion selected from CAB<sup>-</sup>, FAP<sup>-</sup>, FAB<sup>-</sup> and Im<sup>-</sup>,

CAB<sup>-</sup> conforms to formula (II-1)




y1 is 1, 2, 3 or 4,  
 x1 is 0, 1, 2 or 3 and  
 R<sup>0</sup> is alkyl, aryl, fluorinated alkyl, fluorinated aryl, cycloalkyl or alkylaryl, with

**MERCK-3134**

the condition that R<sup>0</sup> may be hydrogen if y<sub>1</sub> is >2,

FAP<sup>-</sup> conforms to formula (II-2)




p<sub>2</sub> is 1 to 20,

m<sub>2</sub> is 0, 1, 2 or 3,

y<sub>2</sub> is 1, 2, 3 or 4,

FAB<sup>-</sup> conforms to formula (II-3)




p<sub>3</sub> is 1 to 20,

m<sub>3</sub> is 0, 1, 2 or 3,

y<sub>3</sub> is 1, 2, 3 or 4,

Im<sup>-</sup> conforms to formula (II-4)



X is carbon or sulfur,

p<sub>4</sub> is 0 to 20 and  $0 \leq m_4 \leq 2p_4+1$ ,

q is 0 to 20 and  $0 \leq k \leq 2q+1$ ,

y<sub>4</sub> is 1 or 2,

m<sub>4</sub> is 0 if p<sub>4</sub> is 0, and

k is 0 if q is 0,

where the carbon atoms of the alkyl chain of the formula II-4 may be bonded to one another by single bonds, and the resultant alkylene chain may in turn be partially or fully substituted by F,

said process comprising reacting a compound of formula XXIX



where R' and Y<sup>-</sup> has one of the meaning indicated in the case of formula XXVIII,

with an aromatic cyclic or heterocyclic compound R''.

**MERCK-3134**

37. (Withdrawn): A compound according to formula XXIX




in which

R' is aryl or heteroaryl,

Y<sup>-</sup> is an anion selected from CAB<sup>-</sup>, FAP<sup>-</sup>, FAB<sup>-</sup> and Im<sup>-</sup>,

CAB<sup>-</sup> conforms to formula (II-1)




y1 is 1, 2, 3 or 4,

x1 is 0, 1, 2 or 3,

R<sup>0</sup> is alkyl, aryl, fluorinated alkyl, fluorinated aryl, cycloalkyl or alkylaryl, with the condition that R<sup>0</sup> may be hydrogen if y1 is >2,

FAP<sup>-</sup> conforms to formula (II-2)




p2 is 1 to 20,

m2 is 0, 1, 2 or 3,

y2 is 1, 2, 3 or 4,

FAB<sup>-</sup> conforms to formula (II-3)




p3 is 1 to 20,

m3 is 0, 1, 2 or 3,

y3 is 1, 2, 3 or 4,

Im<sup>-</sup> conforms to formula (II-4)



X is carbon or sulfur,

p4 is 0 to 20 and  $0 \leq m4 \leq 2p4+1$ ,

q is 0 to 20 and  $0 \leq k \leq 2q+1$ ,

**MERCK-3134**



y<sub>4</sub> is 1 or 2,  
m<sub>4</sub> is 0 if p<sub>4</sub> is 0, and  
k is 0 if q is 0,

where the carbon atoms of the alkyl chain of the formulae II-4 may be bonded to one another by single bonds, and wherein the resultant alkylene chain may in turn be partially or fully substituted by F;

with the provisos that

if X is sulfur, y<sub>4</sub> is 2, and if X is carbon, y<sub>4</sub> is 1 and p<sub>4</sub> or q ≥ 1.

38. (Withdrawn; Currently Amended): In a method of colouring plastics and plastic fibres, preparing for the preparation of flexographic printing inks, ball-point pen pastes, or stamp ink, coloring ~~colouring~~ leather and paper, preparing cosmetic formulations, or coloring in biochemistry, biology, medicine, analytics or electronics, the improvement wherein a dye according to claim 1 is used for coloring.

39. (Withdrawn): In a method of using a dye in data acquisition systems, reprography, in ink microfilters, in photogalvanics, laser technology or the photo industry, the improvement wherein said dye is a dye according to claim 1.

40. (Withdrawn): In a method of using a dye for CD recorders, DVD recorders (DVD+R, DVD+RW), Blu-ray disc (BD-ROM, BD-R, BD-RE), computer to plate, laser filters, laser marking or photopolymerisation, the improvement wherein said dye is a dye according to claim 1.

41. (Previously Presented): A dye according to Claim 28, wherein CAT<sup>+</sup> is a cation of a polymethine dye.

42. (Previously Presented): A dye according to Claim 28, wherein p<sub>2</sub> is 1, 2, 3, 4, 5, 6, 7 or 8.

**MERCK-3134**

43. (Previously Presented): A dye according to Claim 28, wherein p2 is 2, 3 or 4.

44. (Previously Presented): A dye according to Claim 28, wherein Y<sup>-</sup> is PF<sub>3</sub>(C<sub>2</sub>F<sub>5</sub>)<sub>3</sub>, PF<sub>3</sub>(C<sub>4</sub>F<sub>9</sub>)<sub>3</sub>, PF<sub>3</sub>(C<sub>3</sub>F<sub>7</sub>)<sub>3</sub> or PF<sub>4</sub>(C<sub>2</sub>F<sub>5</sub>)<sub>2</sub>.